



Rowlandson, J., Woodman, T., Tennison, S., Edler, K., & Ting, V. P. (2019). Influence of Aromatic Structure on the Thermal Behaviour of Lignin. *Waste and Biomass Valorization*.
<https://doi.org/10.1007/s12649-018-0537-x>

Peer reviewed version

Link to published version (if available):
[10.1007/s12649-018-0537-x](https://doi.org/10.1007/s12649-018-0537-x)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Influence of Aromatic Structure on the Thermal Behaviour of Lignin

Jemma L. Rowlandson^a, Timothy J. Woodman^b, Steve R. Tennison^c,
Karen J. Edler^d and Valeska Ting^{*a}

This file contains:

Supporting Table S1

Supporting Figures S1- S4

-
- [a] Department of Mechanical Engineering
University of Bristol
Bristol, BS8 1TR
*E-mail: V.Ting @bristol.ac.uk
- [b] Department of Pharmacy & Pharmacology,
University of Bath
Bath, BA2 7AY
- [c] Carbon Tex Ltd.
Chancery House
30 St. Johns Road
Woking GU21 7SA
- [d] Department of Chemistry
University of Bath
Bath, BA2 7AY

Supporting information for this article is given via a link at the end of the document.




Figure S1- FTIR spectra of (—) ground biomass (hemp herds, rice husk or eucalyptus chips), the resulting (---)pulp, and (coloured line) lignin following organosolv extraction. Spectra were recorded from 4000 to 600 cm⁻¹, over 32 scans with a resolution of 4 cm⁻¹. Spectra have been offset along the transmission axis, and the region between 1800 and 600 cm⁻¹ has been focused on for clarity. Full spectra of flax and other biomass investigated is included in the supplementary information.

Table S1- Assignment of main lignin 13 C 1 H cross-peaks in the HSQC spectra shown in Figure 3 and 4.

Label	Typical peak (δ_C/δ_H)	Lignin type				Assignment
		Rice husk	Flax	Hemp	Eucalyptus	
B $_{\beta}$	53.6/3.45	53.9/3.46	53.8/3.47	53.9/3.48	53.6/3.49	C $_{\beta}$ -H $_{\beta}$ in phenylcoumaran substructures
C $_{\beta}$	54.2/3.05	54.3/3.06	54.2/3.07	54.2/3.06	54.3/3.07	C $_{\beta}$ -H $_{\beta}$ in resinol substructures
OCH $_3$	56.3/3.75	56.1/3.76	56.1/3.76	56.3/3.75	56.3/3.75	C-H in methoxyls
A $_{\gamma}$	60.3/3.63	60.3/3.60	60.4/3.60	60.4/6.63	60.4/3.65	β -O-4 substructures
I $_{\gamma}$	61.1/4.05	60.4/4.05	60.3/4.04	60.4/4.05	60.3/4.05	C $_{\gamma}$ -H $_{\gamma}$ in <i>p</i> -hydroxycinnamyl alcohol end groups
A' & A''	63.8/3.83-4.30	n/a	n/a	n/a	n/a	C $_{\gamma}$ -H $_{\gamma}$ in γ -acylated β -O-4' substructures
C $_{\gamma}$	63.6/3.63	63.3/3.71	63.3/3.71	63.3/3.71	63.3/3.72	C $_{\gamma}$ -H $_{\gamma}$ in phenylcoumarane substructures
B $_{\gamma}$	71.3/3.77 & 4.09	71.3/3.75 & 4.05	71.3/3.75 & 4.05	71.7/3.80 & 4.05	71.7/3.80 & 4.06	C $_{\gamma}$ -H $_{\gamma}$ in resinol substructures
A $_{\beta}$	80.4/4.5. 84.4/4.4 & 85.6	84.2/4.31	84.2/4.31 & 86.4/4.12	84.3/4.32 & 86.5/4.13	84.1/4.32 & 86.5/4.13	C $_{\beta}$ -H $_{\beta}$ in β -O-4' substructures (A)
A $_{\alpha}$	71.9/4.80	71.7/4.76	71.7/4.77	71.8/4.77	72.6/4.89	C $_{\alpha}$ -H $_{\alpha}$ in β -O-4' substructures (A) and γ -acetylated β -O-4' substructures (A' & A'')
D $_{\alpha}$	82.1/5.12	n/a	n/a	n/a	77.8/5.12	C $_{\alpha}$ -H $_{\alpha}$ in spirodienone substructures
A $_{\beta}$ (G)	84.1/4.29	84.2/4.31	84.2/4.31	84.3/4.32	84.1/4.32	acylated β -O-4 substructures
B $_{\alpha}$	85.6/4.62	85.5/4.62	85.6/4.64	85.7/4.64	85.8/4.64	C $_{\alpha}$ -H $_{\alpha}$ in resinol substructures
A $_{\beta}$ (S)	86.5/4.11	n/a	86.4/4.12	86.5/4.13	86.5/4.13	C $_{\beta}$ -H $_{\beta}$ in β -O-4' substructures (A)
C $_{\alpha}$	87.0/5.43	87.6/5.45	87.6/5.45	87.6/5.45	87.6/5.44	C $_{\alpha}$ -H $_{\alpha}$ in phenylcoumarane substructures
S $_{2,6}$	104.2/6.63	104.1/6.66	104.1/6.66	104.1/6.66	104.1/6.63	C $_{2,6}$ - H $_{2,6}$ in syringyl unit
S' $_{2,6}$	106.7/7.36 & 7.21	104.7/7.35	n/a	106.8/7.33	106.8/7.33	C $_{2,6}$ - H $_{2,6}$ in oxidised (C $_{\alpha}$ =O) syringyl unit
G $_2$	110.5/6.90	110.8/6.93	110.8/6.93	110.9/6.93	111.0/6.93	C $_2$ -H $_2$ in guaiacyl unit
FA $_2$	111.8/7.34	111.7/7.33	112.4/7.33	n/a	110.7/7.48	Ferulic acid
G $_5$	115.2/6.71 & 6.94	115.7/6.77 & 6.93	115.7/6.76 & 6.94	115.7/6.78 & 6.95	115.8/6.76 & 6.95	C $_5$ -H $_5$ in guaiacyl unit
G $_6$	119.5/6.83	119.1/6.77	119.1/6.78	119.3/6.77	119.4/6.78	C $_6$ -H $_6$ in guaiacyl unit
FA $_6$	123.1/7.1	121.6/7.06	n/a	n/a	n/a	Ferulic acid
J $_{\beta}$	126.4/6.94	126.2/6.96	126.1/6.97	126.2/6.96	126.7/7.01	Cinnamaldehyde end group
H $_{2,6}$	128.1/7.27	128.3/7.24	n/a	128.3/7.24	n/a	C $_{2,6}$ - H $_{2,6}$ in <i>p</i> -hydroxyphenyl units
I $_{\alpha}$	130.6/6.3	130.7/6.32	130.8/6.32	130.8/6.33	130.9/6.32	C $_{\alpha}$ -H $_{\alpha}$ in cinnamyl alcohol end-groups
PCA $_{2,6}$	130.7/7.48	130.6/7.47	n/a	130.7/7.51	n/a	C $_{2,6}$ -H $_{2,6}$ in <i>p</i> -coumaric acid



Figure S2- Images of industrial organosolv lignin derived chars carbonised from room temperature to (A) 350 °C and (B) 900 °C, at 5 °C/min, held at temperature for 30 minutes, under a flow of argon (500 ml/min)




Figure S3- Images of industrial organosolv lignin samples heat in a conventional oven to the temperatures listed. By 150 °C lignin had formed a melt. Expansion of the sample onset between 150 and 200 °C, and continued to the maximum temperature investigated of 250 °C. On removal from oven the 250 °C char deflated.




Figure S4- Nitrogen sorption isotherm for char carbonised under argon from industrial organosolv lignin at 900 °C for 30 mins.